

## The XMM-LSS survey. Survey design and first results

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

JCAP09(2004)011

(<http://iopscience.iop.org/1475-7516/2004/09/011>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 159.149.46.61

The article was downloaded on 20/02/2012 at 15:58

Please note that [terms and conditions apply](#).

# The XMM-LSS survey. Survey design and first results\*

Marguerite Pierre<sup>1</sup>, Ivan Valtchanov<sup>1,19</sup>, Bruno Altieri<sup>2</sup>, Stefano Andreon<sup>3</sup>, Micol Bolzonella<sup>4,20</sup>, Malcolm Bremer<sup>5</sup>, Ludovic Disseau<sup>6,7</sup>, Sergio Dos Santos<sup>1</sup>, Poshak Gandhi<sup>7</sup>, Christophe Jean<sup>8</sup>, Florian Pacaud<sup>1</sup>, Andrew Read<sup>9,21</sup>, Alexandre Refregier<sup>1</sup>, Jon Willis<sup>7,10,22</sup>, Christophe Adami<sup>11</sup>, Danielle Alloin<sup>7</sup>, Mark Birkinshaw<sup>5</sup>, Lucio Chiappetti<sup>4</sup>, Aaron Cohen<sup>12</sup>, Alain Detal<sup>8</sup>, Pierre-Alain Duc<sup>1</sup>, Eric Gosset<sup>8</sup>, Jens Hjorth<sup>13</sup>, Laurence Jones<sup>9</sup>, Olivier Le Fevre<sup>11</sup>, Carol Lonsdale<sup>14</sup>, Dario Maccagni<sup>4</sup>, Alain Mazure<sup>11</sup>, Brian McBreen<sup>15</sup>, Henry McCracken<sup>11,23</sup>, Yannick Mellier<sup>16</sup>, Trevor Ponman<sup>9</sup>, Hernan Quintana<sup>10</sup>, Huub Rottgering<sup>17</sup>, Alain Smette<sup>8</sup>, Jean Surdej<sup>8</sup>, Jean-Luc Starck<sup>1</sup>, Laurent Vigroux<sup>1</sup> and Simon White<sup>18</sup>

<sup>1</sup> CEA/DSM/DAPNIA, Service d'Astrophysique, Saclay, F-91191 Gif sur Yvette, France

<sup>2</sup> XMM Science Operations Centre, Villafranca, Spain

<sup>3</sup> Osservatorio Astronomico di Brera, Milano, Italy

<sup>4</sup> IASF, Milano, Italy

<sup>5</sup> Department of Physics, University of Bristol, UK

<sup>6</sup> Ecole Normal Supérieure, Paris, France

<sup>7</sup> European Southern Observatory, Santiago, Chile

<sup>8</sup> Institut d'Astrophysique et de Géophysique, Université de Liège, Belgium

<sup>9</sup> School of Physics and Astronomy, University of Birmingham, UK

<sup>10</sup> Pontificia Universidad Católica, Santiago, Chile

<sup>11</sup> Laboratoire d'Astrophysique, Marseille, France

<sup>12</sup> Naval Research Laboratory, Washington, USA

<sup>13</sup> Astronomical Observatory, Copenhagen, Denmark

<sup>14</sup> Infrared Processing and Analysis Center, Caltech, USA

<sup>15</sup> Physics Department, University College, Dublin, Republic of Ireland

<sup>16</sup> Institut d'Astrophysique, Paris, France

<sup>17</sup> Leiden Observatory, Leiden, The Netherlands

<sup>18</sup> Max Planck Institut für Astrophysik, Garching bei München, Germany

E-mail: [mpierre@cea.fr](mailto:mpierre@cea.fr)

\* Paper based on observations obtained with the XMM,CFH, ESO (Prg: P70. A-0283; .A-0733), VLA, CTIO and Las Campanas observatories.

Received 24 June 2004

Accepted 23 August 2004

Published 20 September 2004

Online at [stacks.iop.org/JCAP/2004/i=09/a=011](http://stacks.iop.org/JCAP/2004/i=09/a=011)

doi:10.1088/1475-7516/2004/09/011

**Abstract.** The XMM Large Scale Structure survey (XMM-LSS) is a medium deep large area X-ray survey. Its goal is to extend large scale structure investigations attempted using ROSAT cluster samples to two redshift bins between  $0 < z < 1$  while maintaining the precision of earlier studies. Two main goals have constrained the survey design: the evolutionary study of the cluster–cluster correlation function and of the cluster number density. The adopted observing configuration consists of an equatorial mosaic of 10 ks pointings, separated by  $20'$  and covering  $8^\circ \times 8^\circ$ , giving a pointsource sensitivity of  $\sim 5 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$  in the 0.5–2 keV band. This will yield more than 800 clusters of galaxies and a sample of X-ray AGN with a space density of about  $300 \text{ deg}^{-2}$ . We present the expected cosmological implications of the survey in the context of  $\Lambda$ CDM models and cluster evolution. We give an overview of the first observational results.

The XMM-LSS survey is associated with several other major surveys, ranging from the UV to the radio wavebands, which will provide the necessary resources for X-ray source identification and further statistical studies. In particular, the associated CFHTLS weak lensing and AMiBA Sunyaev–Zel’dovich surveys over the entire XMM-LSS area will provide for the first time a comprehensive study of the mass distribution and of cluster physics in the universe on scales of a few hundred Mpc. We describe the main characteristics of our wavelet-based X-ray pipeline and source identification procedures, including the classification of the cluster candidates by means of a photometric redshift analysis. This permits the selection of suitable targets for spectroscopic follow-up. We present preliminary results from the first 25 XMM-LSS pointings: X-ray source properties, optical counterparts, and highlights from the first Magellan and VLT/FORS2 spectroscopic runs as well as preliminary results from the NIR search for  $z > 1$  clusters. The results are promising and, so far, in accordance with our predictions. In particular: (1) we reproduce the  $\log N$ – $\log S$  distribution for point sources obtained from deeper surveys at our sensitivity; (2) we find a cluster number density of 15–20  $\text{deg}^{-2}$ ; (3) for the first time, we statistically sample the group mass regime at a redshift out to  $\sim 0.5$ .

**Keywords:** dark matter, lensing

<sup>19</sup> Present address: Imperial College, London, UK.

<sup>20</sup> Present address: Osservatorio Astronomico di Bologna, Bologna, Italy.

<sup>21</sup> Present address: University of Leicester, UK.

<sup>22</sup> Present address: University of Victoria, Canada.

<sup>23</sup> Present address: Institut d’Astrophysique, Paris, France.

---

**Contents**

<b>1. Introduction</b>	<b>3</b>
1.1. The role of clusters in cosmology . . . . .	3
1.2. The quest for clusters . . . . .	4
1.3. The power of XMM . . . . .	6
1.4. The goals of the XMM-LSS . . . . .	6
1.5. The XMM-LSS consortium . . . . .	7
<b>2. Survey design and cosmological implications</b>	<b>7</b>
2.1. Defining the survey . . . . .	7
2.2. Constraining cosmology: quantitative assessments . . . . .	8
<b>3. Associated multi-wavelength surveys: an overview</b>	<b>11</b>
<b>4. X-ray source lists</b>	<b>13</b>
4.1. Source detection and statistics . . . . .	14
4.2. Extended sources and visual check . . . . .	14
<b>5. Source classification</b>	<b>16</b>
5.1. Clusters . . . . .	16
5.2. AGNs and QSO . . . . .	18
<b>6. First spectroscopic campaigns</b>	<b>20</b>
6.1. Strategy . . . . .	20
6.2. Overview of the results . . . . .	21
<b>7. First NIR imaging campaign</b>	<b>23</b>
<b>8. Conclusion and prospects</b>	<b>23</b>
8.1. The newly discovered clusters . . . . .	23
8.2. Cluster identification refinements . . . . .	25
8.3. Active galactic nuclei . . . . .	27
8.4. Survey products . . . . .	28
<b>Acknowledgments</b>	<b>28</b>
<b>References</b>	<b>28</b>

---

**1. Introduction****1.1. The role of clusters in cosmology**

We have come a long way since the Virgo cluster of galaxies was discovered as an accumulation of ‘nebulae’ by Messier in 1781, the first time the term ‘cluster’ was used by Shapley and Ames in 1926, and the first inference of large amounts of ‘dark matter’ in the Virgo cluster by Zwicky in 1933, to the point where clusters of galaxies are now routinely used as cosmological tools. This progress is intimately related to the development and success of the standard cosmological model over the past 50 years, following ever-deeper

insights into the early evolution of the Universe and the growing power of numerical simulations. From a simple theoretical point of view, clusters of galaxies—the most massive bound structures in the Universe—are objects having a mass of the order of  $10^{14-16} M_{\odot}$  growing by accretion at a rate governed by the initial density fluctuation spectrum, the cosmological parameters, and the nature and amount of dark matter as well as the nature of the dark energy. Their three-dimensional space distribution and number density as functions of cosmic time constrain cosmological parameters in a unique way. Non-gravitational effects accompanying cluster formation render the picture more complicated, but compared to galaxies, clusters still offer considerable advantages for large scale structure (LSS) studies: they can provide complete samples of objects over a very large volume of space, and they are in crucial respects simpler to understand. The extent (and mass) of clusters can be traced by their X-ray emission while the theory describing their formation (biasing) and evolution from the initial fluctuations can be tested with numerical simulations. Such a level of understanding does not exist for galaxies—which have reached a highly non-linear stage—and there is even less for QSO and AGN formation. The resulting cluster LSS counts can constrain cosmological parameters, independently of cosmic microwave background (CMB) and supernova (SN) studies, since they do not rely on the same physical processes. In addition, they can also be used to test fundamental assumptions of the standard paradigm, such as the gravitational instability scenario. A quantitative overview of the cosmological implications of cluster surveys can be found for instance in [22].


Conversely, given a cosmological model, a large and deep statistical sample of clusters would provide long-awaited information linking cluster physics, non-linear phenomena involved in cluster evolution, and scaling relations. Further, cluster number counts as a function of both the redshift and X-ray luminosity (or any observable) provide strong consistency tests on assumptions made in modelling the mass–luminosity (or mass–observable) relation and cosmological models involving dark energy [24]. This is timely, since following the WMAP<sup>24</sup> results, independent cosmological constraints from both the early and the local universe must be integrated into a consistent framework.

## 1.2. The quest for clusters

Initiated by Abell in 1958 over the whole sky in the optical, the systematic search for clusters underwent a boost of activity in the X-ray waveband during the EINSTEIN (1978–1981) era. The first X-ray rockets and satellites had revealed the existence of X-ray emission associated with clusters. The UHURU (1970) data, in particular, showed that this emission is thermal and originates in a hot diffuse gas trapped in the gravitational potential of the cluster. The Extended EINSTEIN Medium Sensitivity Survey (EMSS<sup>25</sup>) provided some 730 serendipitous X-ray sources extracted from pointed observations down to a sensitivity of  $1.5 \times 10^{-13} \text{ erg cm}^{-2} \text{ s}^{-1}$  in the 0.3–3.5 keV band. A sub-sample of 67 sources identified as clusters of galaxies in the  $0.14 < z < 0.6$  range [20] suggested, for the first time, a mild evolution in the cluster number density. Next, the ROSAT All-Sky-

<sup>24</sup> <http://lambda.gsfc.nasa.gov/product/map/>

<sup>25</sup> <http://xml.gsfc.nasa.gov/archive/catalogs/9/9015/>



**Figure 1.** Overview of existing X-ray cluster surveys as a function of area and flux. The area plotted is the maximum area of each survey; the flux plotted is the flux at which the survey area is half the maximum. The light filled circles indicate surveys covering contiguous area, while the blue circles represent serendipitous surveys; the stars show the position of the survey presented in this paper, the XMM-LSS survey (current stage and the foreseen final coverage). References: MACS [17], RASS1-BS [15], REFLEX/NORAS [7, 8], EMSS [20], BMW [34], Bright SHARC [39], 160 deg<sup>2</sup> [46], NEP [23], WARPS [35], RDCS [41], RIXOS [12], SHARC-S [14], ROX [16]. Other cluster surveys are CIZA [19], XCS [40] (relevant information not available). For eBCS [18], the shape of the flux/area curve was assumed to be similar to that of REFLEX.

Survey<sup>26</sup> (RASS, 1990–1991) provided the first sample of X-ray clusters over the entire sky, and thus a fundamental resource for LSS studies. Cosmological implications have been investigated in detail from the southern REFLEX ( $3 \times 10^{-12}$  erg cm<sup>-2</sup> s<sup>-1</sup> in the 0.2–2.4 keV band) and north ecliptic pole samples (down to  $3 \times 10^{-14}$  erg cm<sup>-2</sup> s<sup>-1</sup>) by Böhringer *et al* [8] and Henry *et al* [23], respectively.

In parallel, following on the EMSS achievement, serendipitous searches for distant clusters in deep pointed ROSAT observations led to the discovery of clusters out to  $z \sim 1.2$  with a modest evolution of the high luminosity cluster population; a summary of the main X-ray cluster surveys is presented in figure 1.

<sup>26</sup> <http://heasarc.gsfc.nasa.gov/docs/rosat/survey/>. In survey mode the PSF HPD is  $\sim 130''$  at 1 keV (from EXSAS command `calc/psf`).

Both structure and luminosity evolution studies are consistent with hierarchical models of structure formation in a flat low density universe with matter density  $\Omega_m \sim 0.3$  and amplitude of mass fluctuations on an  $8h^{-1}$  Mpc scale,  $\sigma_8 \sim 0.7\text{--}0.8$  (see [42], for a review). In parallel, the quest for clusters in the optical made its own way from automated searches on digitized sky survey plates to deep multi-colour CCD imaging. Indeed, beyond  $z \geq 0.8$ , detecting clusters only from galaxy overdensity in a single band is severely hampered by the faint background population, so that the use of fine-tuned photometric redshift information becomes mandatory (see for example section 5.2). However, given the limitations on the accuracy of such methods and various underlying hypotheses about galaxy evolution, this usually yields large numbers of high- $z$  candidates, many of them simply being portions of cosmic filaments seen in projection. One will thus always require, in default of extensive optical spectroscopic campaigns, an ultimate confirmation from the X-ray band, to assess the presence of deep potential wells.

### 1.3. The power of XMM

At the time the XMM project was initiated<sup>27</sup>, the case for surveys was not as compelling as it is today. Ten years after the completion of the RASS, and following considerable steps forward in our knowledge of cluster physics (e.g. [36]), XMM is in a position to open a new era for X-ray surveys. Its high sensitivity, considerably better PSF than the RASS (FWHM  $\sim 6''$  on axis<sup>28</sup>) and large field of view ( $30'$ ) make it a powerful tool for the study of extragalactic LSS. In this respect, two key points may be emphasized. First, a high galactic latitude field observed with XMM at medium sensitivity ( $\sim 0.5\text{--}1 \times 10^{-14}$  erg cm<sup>-2</sup> s<sup>-1</sup>) is ‘clean’ as it contains only two types of objects, namely QSOs (pointlike sources) and clusters (extended sources) well above the confusion limit. Second, if clusters more luminous than  $L_{[2-10]} \sim 3 \times 10^{44} h_{70}^{-2}$  erg s<sup>-1</sup> are present at high redshift, they can be detected as extended sources out to  $z = 2$ , in XMM exposures of only 10 ks. XMM is a powerful wide angle X-ray imager, with a sensitivity to extended sources which will remain unrivalled in the coming decade. In parallel, progress in optical wide field imaging makes it possible to undertake optical identification of faint X-ray sources over tens of square degrees. Consequently, we are now in a position to probe the evolution of the cosmic network traced by clusters and QSOs over large volumes of the Universe to high redshift.

### 1.4. The goals of the XMM-LSS

Each new generation of instruments brings major improvements over its predecessors. While the BCS sample presented hints of structure in the sky distribution of X-ray bright clusters, REFLEX was the first X-ray survey to systematically address LSS with clusters in the nearby universe (450 clusters with  $z \leq 0.2$ ). Ten years later, there is a clear need to investigate the evolution of the cosmic network out to  $z \sim 1$ . In this context, we have designed a survey to yield some 800 clusters in two redshift bins with  $0 < z < 1$ : the XMM Large Scale Structure Survey (XMM-LSS). This simple goal has set the sensitivity and the

<sup>27</sup> First discussions proposing 27 telescopes in 1982; acceptance in 1988.

<sup>28</sup> The XMM PSF broadens and becomes irregular with increasing off-axis angle; for instance, at 1.5 keV, the radius at which 90% of the energy is encircled is  $\sim 50''$  and  $65''$  for off-axis angles of 0 and  $12'$  respectively for the pn detector; HPD is  $15''$  on-axis at 1.5 keV (cf XMM Users Handbook).



coverage of the XMM-LSS and, as shown in section 2.1, implies an X-ray sensitivity about two orders of magnitude deeper than REFLEX. Consequently, this survey will determine how the cluster number density evolves—a hotly debated topic. It will also trace the LSS as defined by X-ray QSOs out to redshifts of  $\sim 4$ . The cosmological implications of the XMM-LSS are summarized in section 2. In addition, the proposed X-ray survey is associated with several other major new surveys (optical, IR, radio, UV). This will provide a new data set that can be used to study the evolution of clusters, cluster galaxies, and star and AGN formation as a function of environment, in the context of structure formation. An overview of these capabilities is presented in section 3. In the following sections we present a summary of results derived from the XMM AO-1 data set, demonstrating the feasibility of the project. Specifically, in section 4, we describe our current X-ray source identification procedure, which led to the first spectroscopic and NIR follow-up campaigns in Autumn 2002 (sections 5 and 6). Future activities and immediate improvements are presented in section 7.

### 1.5. The XMM-LSS consortium

The wide scope of the project has motivated the assembly of a large consortium to facilitate both the data reduction/management and the scientific analysis of the survey. The XMM-LSS Consortium comprises some fifteen European and Chilean institutes. The project is presented in detail on the following website: [http://vela.astro.ulg.ac.be/themes/spatial/xmm/LSS/index\\_e.html](http://vela.astro.ulg.ac.be/themes/spatial/xmm/LSS/index_e.html).

## 2. Survey design and cosmological implications

### 2.1. Defining the survey

Given our driving goal of extending the REFLEX achievements to high redshift (section 1.4), the following objectives drove the design of the survey.

- *Measure the cluster correlation function in two redshift bins spanning  $0 < z < 1$ , with a good level of accuracy.* A first estimate using the Hubble volume simulations<sup>29</sup> of a  $\Lambda$ CDM universe showed that some 400 clusters per redshift bin are necessary in order to obtain an accuracy of 15–20% on the correlation length; this is comparable to the size of the low- $z$  REFLEX sample. Subsequent implications for the accuracy of the determination of the cosmological parameters are summarized in section 2.2.
- *Probe a comoving length which is significantly larger than  $100h^{-1}$  Mpc at  $z \sim 1$ , the characteristic scale in the galaxy power spectrum of the local universe (e.g. [26]).* This constraint corresponds to an opening survey angle of  $\sim 10^\circ$  at  $z = 1$  (i.e.  $400h^{-1}$  Mpc).
- *Find the best compromise between the two above constraints in order to minimize the necessary XMM observing time.* Quantitatively, we used the Press–Schechter formalism [37] and the mass–temperature relation from simulations to predict the counts of clusters and their X-ray properties in several CDM models [38]. We computed the detection efficiency of clusters, using realistic simulations of XMM images, and studied how this differs from a conventional flux limit.

<sup>29</sup> <http://www.mpa-garching.mpg.de/Virgo/hubble.html>




In order to fulfil the first three conditions above, and assuming the current favoured  $\Lambda$ CDM cosmological model, the optimal survey design was found to be an  $8^\circ \times 8^\circ$  area, paved with 10 ks XMM pointings separated by 20 arcmin (i.e. 9 pointings  $\text{deg}^{-2}$ ). The expected ultimate sensitivity is  $\sim 3 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$  for pointlike sources in the 0.5–2 keV band. This is about 1000 and 10 times deeper than the REFLEX [8] and NEP [23] single area surveys respectively, and provides much better angular resolution ( $6''$  versus  $2'$ ). So far, some  $6 \text{ deg}^2$  have been covered by XMM in the region.

- *Find an adequate survey location.* An equatorial field is optimal, as ground-based follow-up resources from both hemispheres may be used. High galactic latitude and the absence of bright X-ray sources (e.g. nearby clusters) are also required. Moreover, the visibility of the field by XMM must be  $\geq 15\%$ . Given this, only one area in the sky turned out to be favourable: a field centred around  $\alpha = 2^{\text{h}}18^{\text{m}}00^{\text{s}}$ ,  $\delta = -7^{\circ}00'00''$  (at  $b = -58^\circ$ ) with neutral hydrogen column  $1.4 \times 10^{20} < N_{\text{H}} (\text{cm}^{-2}) < 3.5 \times 10^{20}$ . Whereas this region appears to be the best compromise between many astronomical and instrumental constraints, it should be noted that it is not optimal for far infrared observations (above  $25 \mu\text{m}$ ) because of cirrus contamination. The X-ray survey is centred on two deeper areas (one and two square degrees) deriving from XMM guaranteed time programmes (GT). Several surveys are located in the same region (section 3) of which an overview map is presented in figure 2.

## 2.2. Constraining cosmology: quantitative assessments


The prospects that the XMM-LSS cluster catalogue offers for determining cosmological parameters have been studied in detail by Refregier *et al* [38]. We recall here the main issues.

- The cluster counts set strong constraints on the value of the  $(\Omega_{\text{m}} - \sigma_8)$  combination. This combination will also provide a consistency check for the  $\Lambda$ CDM model, and a discrimination between this model and the OCDM model.
- The addition of the cluster two-point correlation function provides a constraint on  $\Gamma$ , the shape of the initial density fluctuation power spectrum.
- With the current survey design, the *simultaneous* expected precision on  $\Omega_{\text{m}}$ ,  $\sigma_8$  and  $\Gamma$  is about 15%, 10%, and 35% respectively. It is important to note that the present uncertainties on  $\sigma_8$  globally result in a factor of 2 uncertainty in our predicted cluster numbers of  $\sim 15 \text{ deg}^{-2}$  (i.e. from 600 to 1200 cluster detections expected within  $0 < z < 1$ ). The high sensitivity to  $\sigma_8$  is not surprising, as it is precisely this which makes cluster counts a good measure of this parameter. This uncertainty can be reduced by analysing about  $15 \text{ deg}^2$  of the XMM-LSS, the minimum area required to improve upon the current measurements of  $\sigma_8$  in the presence of shot noise and cosmic variance.
- The sensitivity of the XMM-LSS survey allows the entire cluster population ( $\geq 2 \text{ keV}$ ) to be detected out to a redshift of 0.6, and will unveil the nearby group population (the cluster selection function can be found in [38]). Rather few galaxy groups have been well studied in the nearby universe, whereas they are believed to constitute the majority of the mass, and of the baryon reservoir. They are also the building block



**XMDs & VVDS deep**  
 VVDS *wide*  
**XMM Subaru Deep Survey**  
**Spitzer Legacy : SWIRE**  
 NOAO Deep Survey  
**Galex**

**Figure 2.** A large white square indicating the location of the XMM-LSS survey is overlaid on a map of  $N_{\text{H}}$  ( $1.4 \times 10^{20} < N_{\text{H}}(\text{cm}^{-2}) < 3.5 \times 10^{20}$  for the XMM-LSS field). The survey area surrounds two deep XMM surveys based on guaranteed time: the XMM-SSC/Subaru Deep Survey (80 ks exposures in  $1 \text{ deg}^2$ ) and the XMM Medium Deep Survey (XMDs; 20 ks exposures in  $2 \text{ deg}^2$ ) also corresponding to the VIRMOS-DESCART Deep Survey [*deep*], the latter being a collaboration between several instrumental teams: XMM-OM (Liège), XMM-EPIC (IASF-MILANO), XMM-SSC (Saclay); CFHTLS (Saclay, IAP); VIRMOS (LAM, IASF-MILANO, OAB). Also indicated are the positions of the associated DESCART-VIRMOS Deep Survey [*wide*], the SWIRE SPITZER Legacy Survey, the Galex survey and the NOAO deep survey (section 3); the  $8.75 \text{ deg}^2$  UKIDSS survey is centred on the Subaru Deep Survey and the CTIO R- $z'$  imaging covers a region corresponding to the VVDS [*wide*] and [*deep*] surveys. (Note that the centre of the whole XMM-LSS survey has been shifted by  $2^\circ$  southward from its initial position because of the presence of the variable type M7 star Mira Ceti (02 19 20.8–02 58 39) which can reach  $I = 1$ , and cause difficulties with optical mapping.)



**Figure 3.** *First view of the deep X-ray sky on large scales.* Image obtained combining the first 15 XMM-LSS fields mosaicked in true X-ray colours: red 0.3–1.0 keV, green 1.0–2.5 keV, blue 2.5–10.0 keV. The circles indicate the sources found in the RASS, the brightest one being a star, HD14938. The other source, apparently not present in our XMM data, is from the RASS Faint Source Catalogue with only 7 counts, and thus is probably a spurious detection. The displayed region covers 1.6 deg<sup>2</sup>. This is the first time that such an X-ray depth has been achieved over such an area. The improvement with respect to the RASS is striking, with a source density of the order of  $\sim 300$  deg<sup>-2</sup> in the 0.5–2 keV band. The wealth of sources includes supersoft and very hard sources, as well as sources with a wide range of intrinsic extents, giving an indication of the scientific potential of the XMM-LSS survey.

of richer clusters. With increasing redshift, the XMM-LSS is less sensitive to low mass systems. Consequently, the low- $z$  and high- $z$  samples to be used for the study of the LSS will pertain to different cluster mass ranges. But as outlined above, this does not prevent the derivation of strong cosmological constraints. The XMM-LSS high- $z$  clustering properties will be compared to the REFLEX ones, which accurately sample the most massive nearby clusters. Moreover, given the large volume sampled at high redshift, the XMM-LSS is well suited to constraining the abundance of distant massive clusters. For instance, the probability of finding a coma-like cluster in the  $1.5 < z < 2$  range over 64 deg<sup>2</sup> is of the order of  $6 \times 10^{-7}$  for a  $\Lambda$ CDM universe. Finding a few such objects would thus put the current favoured cosmological model into severe troubles, or strongly question our understanding of cluster physics at high redshift.

JCAP09(2004)011

**Table 1.** XMM-LSS X-ray and associated surveys.

Observatory/instrument	(Planned) coverage	Band	Final sensitivity
XMM/EPIC	64 deg <sup>2</sup>	0.2–10 keV	$\sim 5 - 3 \times 10^{-15}$ erg cm <sup>-2</sup> s <sup>-1</sup> <sup>b</sup>
CFHT/CFH12K (VVDS Deep) <sup>a</sup>	2 deg <sup>2</sup> GT	B, V, R, I	26.5, 26.0, 26.0, 25.4 <sup>c</sup>
CFHT/CFH12K (VVDS Wide) <sup>a</sup>	3 deg <sup>2</sup> GO	V, R, I	25.4, 25.4, 24.8 <sup>c</sup>
CFHT/MegaCam	72 deg <sup>2</sup>	u*, g', r', i', z'	25.5, 26.8, 26.0, 25.3, 24.3 <sup>d</sup>
CTIO 4 m/Mosaic	$\sim 16$ deg <sup>2</sup>	R, z'	25, 23.5 <sup>e</sup>
UKIRT/WFCAM	8.75 deg <sup>2</sup>	J, H, K	22.5, 22.0, 21.0 <sup>f</sup>
VLA/A-array <sup>a</sup>	110 deg <sup>2</sup>	74 MHz	275 mJy/beam <sup>g</sup>
VLA/A-array	5.6 deg <sup>2</sup>	325 MHz	4 mJy/beam <sup>h</sup>
OCRA	All XMM-LSS clusters	30 GHz	100 $\mu$ Jy <sup>i</sup>
AMiBA	70 deg <sup>2</sup>	95 GHz	3.0 mJy <sup>j</sup>
SPITZER/IRAC (SWIRE Legacy) <sup>a</sup>	8.7 deg <sup>2</sup>	3.6, 4.5, 5.8, 8.0 $\mu$ m	7.3, 9.7, 27.5, 32.5 $\mu$ Jy <sup>k</sup>
SPITZER/MIPS (SWIRE Legacy) <sup>a</sup>	8.9 deg <sup>2</sup>	24, 70, 160 $\mu$ m	0.45, 6.3, 60 mJy <sup>l</sup>
Galex	$\sim 20$ deg <sup>2</sup>	1305–3000 Å	$\sim 25.5^m$

<sup>a</sup> Complete.<sup>b</sup> For pointlike sources in 0.5–2 keV.<sup>c</sup>  $AB_{\text{Mag}}$ , 5'' aperture.<sup>d</sup>  $S/N = 5$  in 1.15'' aperture.<sup>e</sup> 4  $\sigma$  in 3'' aperture.<sup>f</sup>  $Vega_{\text{Mag}}$ .<sup>g</sup> 30'' resolution; deeper observations planned.<sup>h</sup> 6.3'' resolution.<sup>i</sup> 5 $\sigma$ , detection limit.<sup>j</sup> 6 $\sigma$ , detection limit.<sup>k</sup> 5 $\sigma$ .<sup>l</sup> 5 $\sigma$ .<sup>m</sup>  $AB_{\text{Mag}}$ .

### 3. Associated multi-wavelength surveys: an overview

While optical information remains the primary data base for X-ray source identification, contribution from other wave bands may be critical, for example in the far infrared domain, where many heavily absorbed X-ray QSO, not visible in the optical, are expected to show up. Beyond the necessary identification step, multi-wavelength information provides an overview of the energy emission and absorption processes in astronomical objects, which is vital to our understanding of their physics, formation and evolution. The contiguous design of the XMM-LSS, optimized for large scale studies, provides considerable advantages for complementary observations, compared to serendipitous fields, and the project has developed numerous collaborations at other wavelengths. The main

characteristics of associated surveys are summarized in table 1 and their main science applications are outlined below.

*Optical, NIR and UV imaging.* The imaging of the  $8 \times 8 \text{ deg}^2$  XMM-LSS area is one of the priorities of the Canada–France–Hawaii Legacy Survey<sup>30</sup> (CFHTLS). It is being performed by MegaCam, the one degree field imager built by CEA and installed at the new CFHT prime focus. Imaging of the XMM-LSS region started by mid 2003. The CFHTLS will provide a deep high quality optical multi-colour imaging counterpart of the X-ray sources at a rate of  $25 \text{ deg}^2 \text{ yr}^{-1}$  in at least three colours. Data pipelines and processing have been developed by the TERAPIX<sup>31</sup> consortium which provides object catalogues and astrometric positions for the entire surveyed region. Currently, the optical data used for the identification work, and presented in this paper, mostly pertain to the CFH12K VIRMOS-DESCART VLT Deep Surveys (VVDS [*deep*] and [*wide*], figure 2 and table 1, [27]). An optical cluster catalogue is under construction using both spatial clustering analysis and multi-colour matched filter techniques, in addition to photometric redshift estimates. Moreover, the MegaCam data will form the basis of a weak lensing analysis<sup>32</sup>, whose cosmological constraints will be compared to those provided by the X-ray data on the same region. This will be the first coherent study of LSS on such scales. R +z' images taken by us at CTIO are also being analysed, forming the basis of an independent cluster catalogue (section 5.1, [2]). In addition, deep NIR VLT imaging (J&K) of  $z > 1$  cluster candidates found in the XMM-LSS is performed as a confirmation prior to spectroscopy. Finally, a sub-area of  $8.75 \text{ deg}^2$  of the XMM-LSS field is a high priority target of the Deep Extragalactic Survey part of the UKIRT Deep Sky Survey (UKIDSS<sup>33</sup>). In the UV domain, the XMM-LSS field is one of the targets of the Galax<sup>34</sup> Deep Survey, whose main goal is to map the global history of star formation out to  $z \sim 2$ .

*Spectroscopy.* The standard spectroscopic follow-up is designed to perform redshift measurements for all identified  $0 < z < 1$  X-ray clusters in multi-object-spectroscopy mode, using 4 m and 8 m class telescopes. Current identification procedures and first results are described in later sections. We shall subsequently undertake programmes of advanced spectroscopy that will focus on individual objects, and include high resolution spectroscopy, the measurement of cluster velocity dispersions and QSO absorption line surveys, as well as NIR spectroscopy of our  $z > 1$  cluster candidates.

*Radio.* In the radio waveband, the complete survey region is being mapped using the VLA at 74 and 325 MHz. First results of this low frequency coverage are described by [13]. Radio observations are not only particularly relevant for tracing merger events triggered by structure formation, but also as a useful indicator of galactic nuclear or star-formation activity.

*Sunyaev–Zel’dovich.* Observations (SZ) are also planned. Clusters in the XMM-LSS field are the targets of the prototype OCRA (one-centimetre radiometer array, [10]) instrument. The full XMM-LSS field will be mapped by the OCRA (starting in 2005), and will be an early target of the array for microwave background anisotropy (AMiBA) [29].

<sup>30</sup> <http://cdsweb.u-strasbg.fr:2001/Science/CFHLS/cfhtlscfhtoverview.html>

<sup>31</sup> <http://terapix.iap.fr>

<sup>32</sup> <http://www2.iap.fr/LaboEtActivites/ThemesRecherche/Lentilles/LentillesTop.html>

<sup>33</sup> <http://www.ukidss.org/>

<sup>34</sup> <http://www.srl.caltech.edu/galextech/galex.htm>



This will enable not only the measurement of the Hubble constant, but also a statistical analysis of the physics of the ICM as a function of redshift. For instance, the AMiBA deep survey will be sensitive to clusters with  $M \geq 1.5 \times 10^{14}$  out to  $z < 2$  [28]. A cross-correlation between the AMiBA and XMM maps in the survey region will thus allow us to study the gas properties of X-ray selected clusters and to detect clusters beyond the X-ray detection threshold. Such measurements are complementary to the X-ray and weak lensing surveys, connecting the mass distribution of clusters to the structure of the hot gas they contain. The three data sets together will also provide a direct and independent check of the extragalactic distance scale.

*Infrared.* In the infrared, the SWIRE<sup>35</sup> SPITZER Legacy Programme covered  $\sim 9 \text{ deg}^2$  of the XMM-LSS in seven MIR and FIR wavebands from 4 to 160  $\mu\text{m}$  [30]. The estimated IR source densities per square degree in this area are around 1100/400/130 and 670/150/130 for starbursts/spiral-irregular/AGN in the  $0 < z < 1$  and  $1 < z < 2$  redshift intervals, respectively [48]. This represents a unique X-ray/IR combination in depth and scales to be probed. The coordinated SWIRE/XMM-LSS observations will clarify an important aspect of environmental studies, namely how star formation in cluster galaxies depends on distance from the cluster centre, on the strength of the gravitational potential, and on the density of the ICM. In this respect the XMM-LSS represents the optimum SWIRE field, where galaxy environment, deep NIR imaging and optical spectroscopic properties will be the main parameters in modelling the MIR/FIR activity. Here also, the location of IR AGNs within the cosmic web will help establish their nature. The FIR/X/optical/radio association will also provide valuable insights into the physics of heavily obscured objects, as well as the first coherent study of biasing mechanisms as a function of scale and cosmic time, for hot (XMM), dark (weak lensing), luminous galactic (optical/NIR) and obscured (SWIRE) material.

In summary, we may underline the complementarity of the two major categories of associated surveys. Those covering an area of the order of 10–20  $\text{deg}^2$ , focusing on source environment, will provide new insights on the physics of AGN and clusters. Surveys planned over a  $\sim 70 \text{ deg}^2$  area (MegaCam and AMiBA), in addition to proper LSS studies, are intended as strong consistency tests on assumptions made in modelling the cluster mass/luminosity and mass/temperature relations and cosmological models involving dark energy.

#### 4. X-ray source lists

In addition to the XMDS Guaranteed Time survey consisting of 18 pointings (figure 2), 33 Guest Observer XMM-LSS pointings have been allocated in the first two XMM AOs. The results presented in this section pertain to the first 27 GO XMM-LSS pointings allocated in AO1. Most of the X-ray observations were performed in good background conditions. However, two pointed GO observations (namely B13 and B17) are heavily contaminated by solar flares, having a mean total count-rate approximately 100 times higher than the quiescent background. Effective exposure times range from 5 to 16 ks (after removal of bad time intervals, mainly due to flares). The mean exposure time weighted over the area is about 12 and 9 ks for the MOS and pn detectors respectively, which is close to the nominal survey exposure time of 10 ks. Statistics presented in this section are obtained

<sup>35</sup> <http://www.ipac.caltech.edu/SWIRE>

using the 25 good pointed observations that were used out to an off-axis angle of 10' and cover a total area of 2.18 deg<sup>2</sup>.

#### 4.1. Source detection and statistics

The XMM-LSS pipeline is based on a 3 stage filtering/detection/measurement process initiated by [44]. After applying the standard XMM pre-reduction procedure<sup>36</sup>, photon images are generated in several energy bands, and the 3 detector images are co-added in each band.


This mosaic is subsequently filtered by a multi-resolution algorithm (in counts, to preserve Poisson statistics): a scale-dependent wavelet coefficient threshold, (corresponding to a probability for each event of being a random background fluctuation lower than  $10^{-3}$ ) is computed in wavelet space by histogram auto-convolution, and followed by an iterative reconstruction of the image (MR1, see [43]). The source extraction procedure SExtractor [6] is then applied on the resulting image to obtain a preliminary list of sources. At this stage, extraction parameters are set in order to avoid missing faint sources. Co-adding images allows us to increase the S/N before filtering and, consequently, to lower the detection thresholds in the first step. Third step consists in examining the likelihood of the detections and characterizing their extent by means of a procedure we have developed (Xamin, [33]). The principle is the following: For each SExtractor source, it performs a maximum likelihood profile fit on two raw photon images (PN image, and co-added MOS images) independently, thus yielding a quantitative assessment of the source, taking into account the PSF variation with energy and off-axis radius, as well as other detector characteristics (vignetting, bad pixels, CCD gaps). Two source models are tested: a PSF model provided by the XMM calibration files, and a spherically symmetric  $\beta$ -model (with fixed  $\beta = 2/3$ ) convolved with the PSF. Models are weighted by the exposure maps at the location of the fit. The parameters that are varied for each source are: position, local background, count rates and core radius of the  $\beta$ -model. Position and extension are forced to be the same in each detector, while count rates are adjusted independently for each EPIC detector type. At the end of the analysis Xamin yields a catalogue of some 50 parameters for each source including: source count rates (in each detector type), likelihood of detection, likelihood of source extent, and extension in arcsec. Finally, discrimination between pointlike and extended sources, is performed in the [Detection Likelihood / Extent] parameter space, calibrated by means of extensive image simulations spanning a wide range in source counts and extent. Performing this analysis on the 25 first 10 ks pointings, we detect some 200 sources per square degree down to a flux limit of  $5 \cdot 10^{-15}$  erg cm<sup>-2</sup> s<sup>-1</sup> in the [0.5-2] keV band for a detection likelihood threshold of 20 (assuming a power-law spectrum with a photon index of  $\Gamma = 1.7$ ). Corresponding log  $N$ -log  $S$  curve is shown on figure 5.

#### 4.2. Extended sources and visual check

As clusters of galaxies constitute the core of the project, special care is devoted to the detection and assessment of faint extended sources. The MR1/SExtractor/Xamin hybrid method described above provides a series of parameters used to establish a preliminary

<sup>36</sup> [http://xmm.vilspa.esa.es/external/xmm\\_sw\\_cal/sas\\_frame.shtml](http://xmm.vilspa.esa.es/external/xmm_sw_cal/sas_frame.shtml)





**Figure 4.** This image ( $10' \times 10'$ ) provides an overview of the source types encountered in the survey and of the identification procedure. Overlaid on a 1 h exposure CFH12KI-band image (processed by Terapix) are blue XMM flux contours in 0.5–2 keV, obtained by the MR1 multi-scale wavelet filtering algorithm; the procedure allows the automatic recognition of pointlike sources (green squares); the significance of the lowest wavelet contour is  $10^{-4}$  (equivalent to  $3.7\sigma$  for Gaussian noise, cf [43]). The image is centred on a bright extended X-ray source, corresponding to an obvious nearby cluster (a *NEAR* candidate) for which we measured a redshift of 0.33. Another—much more distant—cluster is apparent in the upper left corner (a *MID* candidate); this object was found to have a redshift of 0.84. Top middle, another extended source without a clear optical counterpart, typically a *DISTANT* candidate. The density of pointlike sources is high, some of them having an obvious optical counterpart, others, none. The red squares and green circles indicate VLA NVSS and 325 MHz radio sources respectively. In the lower right corner, there is a conspicuous double lobe radio galaxy, which is also an X-ray source.

list of extended sources. Currently, the procedure is performed on the 0.5–2 keV catalogue which offers optimal sensitivity considering the observed cluster emission spectra over the  $0 < z < 1$  range, the various components of the background (galactic, particle, solar flares), and galactic absorption. We have developed an automated interface to produce X-ray/optical overlays for every extended source candidate (figure 4). At the end of the


pipeline procedure, each overlay is inspected by eye in order to catch possible instrumental artefacts which could have escaped the pipeline rejection algorithm. Finally, a list of plausible extended sources is issued with an indication of whether they correspond to an obvious overdensity of galaxies. We currently find an extended source density of about  $15\text{--}20 \text{ deg}^{-2}$ , with conspicuous optical cluster (or group) counterparts on the CFH12K I images. At this stage, given the small number of spectroscopically confirmed clusters, it is not possible to make firm statements about these results. The fact that we tend to find somewhat more extended objects than formally predicted would suggest that either we sample the cluster population below the 2 keV limit that was set in our theoretical predictions, using a given set of M–T–L( $z$ ) relations, or that we are more sensitive than assumed. An example of the sensitivity performances for extended sources is displayed in figure 6. This suggests that we reach a sensitivity of at least  $10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$  in the 0.5–2 keV band to extended sources over a large fraction of the detector. This is in agreement with the predictions of Refregier *et al* [38], showing that the cluster selection function is close to a flux limit of  $10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$  for  $z < 1$  clusters. The detailed study of the  $\log N\text{--}\log S$  distribution of extended sources and of their selection function will be the subject of a forthcoming paper.

## 5. Source classification

### 5.1. Clusters

The depths of both the XMM-LSS and of the CFHTLS and CTIO data have been tuned to sample most of the cluster population out to a redshift of 1 (section 2.1 and table 1). Beyond this, massive clusters can still be detected in the X-ray band [44]. A cluster located at  $z \sim 2$  and having a temperature of 7 keV will show up with an apparent temperature of about 2.5 keV, which falls in the most sensitive part of the XMM response. Hence, for such a cluster, the X-ray K-correction is only  $\sim 0.7$  [25]. In contrast,  $z > 1$  clusters do not appear as significant overdensities of galaxies in the optical band. Increasing the depth of the optical survey would not significantly improve the situation, as most of the galaxy light is shifted into the infrared band. We have therefore adopted the following approach and definitions.


- (1) A  $z \leq 1$  cluster candidate is defined as an extended X-ray source corresponding to a significant visual excess of galaxies in the optical wavebands. The multi-colour BVRIz' information is used to enhance the visual contrast of the cluster galaxy density with respect to the background population and to construct optical cluster catalogues following different methods: the Red Cluster Sequence [21], [2], clustering analysis in redshift slices [1], and matched filter [31]. These catalogues are currently used as ancillary information and cross-correlated with the X-ray extended source catalogue. Further, the optical data allow us to assign photometric redshifts to the X-ray cluster candidates in an independent way. First, we apply the public code Hyperz [9] to the entire BVRI galaxy catalogue. For each galaxy, we derive the photometric redshift, corresponding to the minimum  $\chi^2$  computed comparing the observed photometry to the fluxes expected from a set of reference templates (GISSEL98, [11]). Second, we search for the cluster signature by placing a number of apertures in the field of



**Figure 5.** Left: Cumulative  $\log N$ – $\log S$  distribution, in the [0.5–2] keV band, for the sources pertaining to the first 25 XMM-LSS pointings (off-axis  $< 10'$ ,  $N_{\text{counts}} > 20$  and detection likelihood  $> 20$ ). The dashed line is the analytical fit of the deep  $\log N$ – $\log S$  distribution, by [32], the dotted line the determination by [5] for the HELLAS2 survey. Right: Corresponding sky coverage for point sources shows that 90% of the area is covered with a sensitivity better than  $5 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$ . On-going pipeline developments, coupled with extensive image simulations, aim to a deeper completeness limit.

the X-ray detection in order to determine the maximum overdensity in photometric redshift space relative to the ‘field’ distribution. This approach does not assume that the distribution of cluster galaxy members apparent in the optical data is centred on the X-ray detection and it makes no prior assumption of the apparent optical extent of the candidate cluster. The redshift bin in  $N(z_{\text{phot}})$  has been set similar to the precision expected from photometric redshifts, given the set of filters and their depths (typically  $\Delta z = 0.2$ ). Finally, from the region with the largest overdensity detected, we selected the candidate cluster members in different classes of confidence, considering their  $z_{\text{phot}}$ , their error bars and probability functions. The method is illustrated by figures 7 and 8. This procedure enables the pre-selection of the cluster candidates into *NEAR* and *MID* distance classes, corresponding to the  $0 < z < 0.5$  and  $0.5 < z < 1$  ranges respectively, and provides a useful tool for ascribing targets to the various telescopes available for the spectroscopic follow-up.

- (2) A small fraction of the extended sources ( $\leq 1$  per pointing) do not present any significant optical counterpart—occasionally, one or two galaxies visible in the I or  $z'$  band, sometimes nothing at all. These are typically interesting  $z > 1$  cluster candidates




**Figure 6.** The figure shows an XMM source detected as extended with only  $\sim 50$  net counts in the 0.5–2 keV combined image of the three detectors. The overlay on the optical image confirms the presence of a distant cluster ( $z \sim 0.8\text{--}0.9$ ), and even reveals X-ray structure associated with two distinct groups of galaxies. The sources actually lie at an off-axis angle of  $11.5'$ . The collected photons correspond to an emitted flux of  $\sim 1.2 \times 10^{-14}$  or  $\sim 4 \times 10^{-15}$   $\text{erg cm}^{-2} \text{s}^{-1}$ , for a source falling at an off-axis distance of  $11'$  or on-axis respectively, assuming a typical thermal cluster spectrum.

and are classified into the third category, *DISTANT*, to be imaged in the NIR bands. If confirmed, they are the subject of dedicated spectroscopic follow-up programmes.


## 5.2. AGNs and QSO

Active galactic nuclei constitute by far the dominant population of X-ray sources at the XMM-LSS sensitivity. Given the XMM PSF, and consequently the positional accuracy, there is some ambiguity in the identification of pointlike sources, as can be appreciated in figure 4 (note the sources having two or three possible optical counterparts, or for which the optical ID is below the limit of the optical survey). This is a well known drawback, thoroughly studied in deep surveys; but here, the situation is simpler since the XMM-LSS flux limit is well above the confusion limit.





**Figure 7.** Photometric redshift determination around the position of an X-ray cluster candidate (see text). Top: the photometric redshift distribution of the region where the overdensity has been detected (solid line), compared to the reference redshift distribution, obtained from the total catalogue (dotted line). The redshift distributions have been normalized to the respective number of objects. To select the significant overdensities, we plot the Poissonian error bars of the small region  $N(z_{\text{phot}})$ . Bottom: combined photometric redshift probability as a function of redshift of galaxies selected within an aperture of radius  $1'$  whose position within the field maximizes the probability of the redshift ‘peak’. Individual galaxy probabilities are determined from the chi-squared statistic returned from the photometric redshift template fitting procedure. VLT/FORS2 observations showed that this X-ray source indeed corresponds to a structure at  $z \sim 0.9-1$ . See figure 8.



**Figure 8.** The most distant X-ray complex identified so far, showing a clear concentration of galaxies in the  $0.9 < z < 1$  range (green X-ray contours overlaid on a CFH12K VRI composite). Bottom right is the most distant X-ray QSO currently measured in the XMM-LSS at a redshift  $z = 3.3$  (see figure 12); its isophotes are distorted by the immediate vicinity of an X-ray bright emission line galaxy located at  $z = 0.054$ .

## 6. First spectroscopic campaigns

### 6.1. Strategy

The first spectroscopic observations dedicated to the identification and redshift measurement of the XMM-LSS galaxy clusters took place during Autumn 2002. Three nights were allocated at ESO on the VLT/FORS2 instrument and two nights at Las Campanas with the Magellan/LDSS2 instrument. We summarize here the most important outcomes. Galaxy cluster candidates were pre-selected as described above, comprehensive information (X-ray, optical, photometric redshift) being gathered into an HTML database, allowing a rapid overview of the cluster properties. From this, we selected a number of clusters spanning different optical morphology and flux ranges, in order to optimize the allocated spectroscopic time, and to give a representative overview of the XMM-LSS cluster population. For the first spectroscopic campaign, it was decided to conservatively obtain two masks per cluster in order to sample well the cluster galaxies in the central region, and ensure confident redshift determination. Thanks to the high throughput of the

FORS2 instrument in the 0.8–1  $\mu\text{m}$  range, all *MID* galaxy cluster candidates were directed to the VLT sample. The *NEAR* ones were assigned to the Magellan sample, and also as backup sources for VLT, in case of possible poor weather. In order to investigate and quantify the most efficient method for the future, galaxies to be measured were selected using different approaches: simple magnitude cut, colour selection or photometric redshift criteria.


No dedicated spectroscopic runs were planned in 2002 for the QSO and AGN population. However, many pointlike X-ray sources surrounding the selected clusters and present in the corresponding instrumental field of view were included in the spectroscopic masks. For each  $7' \times 7'$  field centred on a galaxy cluster candidate, we first cross-correlated the X-ray pointlike source catalogues with the optical ones obtained from the CFH12K data. We then searched for optical counterparts within a radius of  $5''$  from each X-ray pointlike source and produced catalogues of associations. Using the latter, we overlaid on the optical images the positions of the X-ray sources and their associated optical counterparts, if any. Then, we visually checked on the optical images each X-ray position (no more than ten per field) associated with 0, 1 or more optical counterparts. Depending on the ‘quality’ of the X-ray/optical associations, we sorted the AGN/QSO candidates into different categories characterizing the priority for spectroscopic observations of the optical counterparts. There were three main classes: (1) unambiguous cases where a unique and relatively bright optical source lies within a radius of  $5''$ ; (2) ambiguous or doubtful cases when there were several possible optical counterparts and/or when the optical counterpart(s) was (were) rather faint; (3) rejected cases when there was no, or a much too faint, optical counterpart. The fraction represented by each class was  $\sim 35$ , 25 and 40% respectively. A catalogue containing the first two classes of AGN/QSO candidates was considered for spectroscopic observations of each field. Given the constraints imposed by the cluster spectroscopy (especially, optimal sampling of the cluster cores) and the zones of avoidance within the FORS2 field of view, an average of 2–3 pointlike sources was measured per cluster field. This corresponds to about 30–45 sources  $\text{deg}^{-2}$ .

## 6.2. Overview of the results

Weather and working conditions at the VLT were optimal. In three nights, 12 cluster fields (5 *MID*, 7 *NEAR* –  $z > 0.35$ ) were observed, containing on average about 30 slits per mask, yielding some 700 spectra. In addition, two new nearby compact groups were also observed (with seven and eight galaxies per group), prepared as a backup programme. The overall strategy proved successful. An example spectrum is shown in figure 9. Encouragingly, an estimate of the velocity dispersion (16 cluster galaxy spectra) of a newly discovered 0.85 redshift cluster can be achieved in 2 h, supporting the follow-up of large numbers of distant clusters with relatively little observing time. Seven *NEAR* clusters were observed at Magellan, where one night out of the allocated two suffered from poor meteorological conditions. With an average number of slits of 15 per masks, some 200 spectra were obtained.

The photometric cluster distance estimates proved useful and reliable indicators for the cluster classification (see figure 7, 8). Current results on the measured redshift distribution are shown in figure 10. Given the exploratory selection procedure adopted for this first identification campaign, this cannot be ascribed any cosmological interpretation






**Figure 9.** One of the most distant cluster galaxy measured, at a redshift of  $z = 1.02$  (see figure 8). This good quality spectrum was obtained in 1.5 h with the R600RI + 19 grism on the VLT/FORS2 instrument. The object has magnitudes of  $V = 24.19$ ,  $R = 23.34$ ,  $I = 22.21$  and displays characteristics of an elliptical galaxy having a component of young stars (E + A type).

yet. These measurements, however, enabled us to compute the cluster luminosities and, subsequently, to confirm our sensitivity predictions [38] and to demonstrate that we are sampling a new population of low luminosity (i.e. low mass) objects. Figure 11 displays the luminosity distribution of our current confirmed cluster sample as a function of redshift, compared to the previously known set of X-ray clusters: out to 0.5, we identified a bunch of very faint clusters, thus significantly extending the sample that was inventoried in the very local universe so far ( $z < 0.1$ ). Between  $0.5 < z < 1$ , we are similarly discovering the high redshift counterparts of the known  $z < 0.5$  cluster population. Further statistical analysis of this population will be especially relevant to better constrain the evolution of cluster scaling laws as to the gas properties, since groups are more sensitive than clusters to the interactions with the individual galaxies. From the LSS point of view, low mass groups will provide a precious link between structure underlined by clusters and those by galaxies, thus enlightening the bias controversy. A comprehensive X-ray/optical study of the current *NEAR* and *MID* samples is presented by Willis *et al* [47] and Valtchanov *et al* [45] respectively. The optical properties of the cluster galaxies (luminosity function and colour distribution) are discussed by Andreon *et al* [2].

The optical spectra of the measured X-ray AGNs showed a plethora of properties. The most distant X-ray source measured so far has a redshift of 3.3 (figure 12). The AGN sample is currently under analysis.



**Figure 10.** Redshift distribution of the 21 spectroscopically confirmed clusters and compact groups during the 2002 Magellan/LDSS and VLT/FORS2 runs.

JCAP09(2004)011


## 7. First NIR imaging campaign

In November 2002, we carried out an initial, exploratory observing run with SOFI on the NTT, in order to search for *DISTANT* clusters selected from candidates in the first set of available XMM data. The details will be presented by [3]. Here we note that by selecting faint, extended X-ray sources associated with either blank fields in the CFHT data, or with fields showing a hint of clustering at faint ( $I > 22$ ) magnitude levels in the same data, we were successful at reliably detecting such clusters. Moreover, the X-ray fluxes of these objects are such that if they were placed at higher redshifts they would still be detectable in the XMM-LSS exposures. Thus the survey will enable us to map the cluster distribution to redshifts well above  $z = 1$ . An example is shown in figures 13 and 14.

## 8. Conclusion and prospects

### 8.1. The newly discovered clusters


Following the completion of the AO-1 period, the main outcome of the XMM-LSS project can be summarized as follows: XMM observations of only 10 ks, coupled with a similar time spent on imaging at CFHT or CTIO, can detect a significant fraction of the cluster population out to  $z \sim 1$  (a flux of  $0.8 \times 10^{-14}$  erg cm<sup>2</sup> s<sup>-1</sup> corresponds to a cluster mass of



**Figure 11.** The locus of the XMM-LSS clusters. Clusters observed during the first 2002 spectroscopic campaign are indicated as large filled green circles, while dots indicate all clusters in the literature having X-ray observations (BAX: <http://bax.ast.obs-mip.fr/>). The curved lower edge of the shaded region indicates the expected XMM-LSS ultimate cluster completeness limit ( $\sim 8 \times 10^{-15}$  erg s $^{-1}$  cm $^{-2}$  in 0.5–2 keV), whereas the upper limit is merely to guide the eye (maximum luminosity at which at least one cluster is expected within the full 64 deg $^2$  area, according to  $\Lambda$ CDM model). Although the 2002 exploratory campaign was not intended to be flux limited, the figure well enlightens the tendency of the XMM-LSS to identify significantly lower luminosity (i.e. lower mass) systems than the previous cluster surveys. Luminosity is expressed in the 0.1–2.4 keV band.

$6.7 \times 10^{13} M_{\odot}$  ( $T = 2.7$  keV) at  $z = 1$ , assuming typical scaling laws). Moreover, adding two hours of FORS2 spectroscopic time provides a reliable estimate of cluster velocity dispersion at  $z = 0.85$ . This represents a substantial increase in efficiency compared to former high- $z$  cluster searches.

For the first time we are detecting the numerous population of low luminosity (mass) objects out to  $z \sim 0.5$ ; this will allow a dense mapping of the matter distribution. So far (within  $\sim 4$  deg $^2$ , including the GT area), no massive object has been detected and, following a detailed inspection of the optical images, no giant arcs have been found in the identified clusters. These results are consistent with the Press–Schechter formalism (folded



**Figure 12.** Most distant X-ray AGN detected so far, at a redshift of  $z = 3.3$  (see figure 8); grey bands indicate the position of strong atmospheric lines.


with the current favoured cosmological model) predicting  $2.8$  and  $0.0001$  cluster  $\text{deg}^{-2}$  between  $0 < z < 1$  for clusters more massive than  $10^{14}$  and  $10^{15} M_{\odot}$  respectively, if we assume that only massive clusters are potential strong lenses. Of course, an additional necessary condition is the steepness of the mass profile.

Immediate foreseen improvements are the following. In the next version of the X-ray pipeline, the source flux detection limit will be significantly lowered. This requires further work on the background estimation. In parallel, extensive simulations are being performed to improve the characterization (extent and flux) of the faintest sources as well as to determine the survey selection function. This is complicated by the necessity of taking proper account of the Poisson nature of the data and the fact that the signal to be analysed comes from three different detectors.

## 8.2. Cluster identification refinements

In a second step, we shall systematically investigate overlaps and differences between the optical cluster catalogues and the X-ray extended source catalogue. There are two main reasons why variations should be observed.

- (i) Faint groups or distant clusters may not be unambiguously detected as extended sources because of their low flux or, alternatively, because they host a cooling flow,



**Figure 13.** Extended X-ray source with a very faint optical counterpart in the *I*-band. The NIR image of the field is shown in figure 14.


JCAP09(2004)011

which makes them appear unresolved. Currently, we are confident that any regular object having a typical core radius of the order of  $125 h_{100}^{-1}$  kpc and producing a net number of pn counts of  $\geq 70$  is detected by the pipeline as an extended source [44].

- (ii) There are intrinsic differences between the X-ray and optical catalogues since the methods used to construct them rely on specific assumptions as to galaxy colour and evolution, or on the properties of the IGM at high redshift.

Understanding the discrepancies will not only increase the efficiency of our cluster finding procedure, but also shed light on the much debated topic of cluster formation and evolution. This will also contribute to the improvement of our photometric redshift determination procedure. Finally, output from the weak lensing analysis will reveal large mass concentrations, optimally within the  $0.1 < z < 0.5$  range. Again, the comparison between the X-ray and optical catalogues will be most instructive. Given the current understanding of structure formation, it is difficult to devise a physical process that would prevent gas being trapped or heated within large concentrations of dark matter. However, the surveyed volume is ideally suited to systematically search for ‘dark clump’.





**Figure 14.** J-K colour image of the *DISTANT* cluster candidate flagged in figure 13. This presents a significant overdensity of red objects. The brightest cluster member has  $K_s \sim 17.5 \pm 0.3$ , and the other objects within  $20''$  are generally fainter than  $K_s \sim 18.5$ . These properties are consistent with a cluster at  $z = 1$  or above.

JCAP09(2004)011

### 8.3. Active galactic nuclei

Due to the primary goal of the spectroscopic observations performed so far (confirmation and redshift determination of X-ray selected galaxy clusters), few AGN/QSO candidates could be observed. For this reason, a spectroscopic survey of a unique sample of X-ray selected AGN/QSOs within a large contiguous area of some  $10 \text{ deg}^2$ , typically complete down to  $F_{2-10 \text{ keV}} \sim 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$ ,  $F_{0.5-2 \text{ keV}} \sim 5 \times 10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$  and  $I_{AB} = 22.5$ , is foreseen. Because of the large surface density of AGN/QSO candidates detected in the XMM-LSS field (typically  $>200 \text{ deg}^{-2}$  with  $0 < z < 1$ ), compared to previous surveys, it should be more sensitive to contrasts between voids and peaks than previous surveys, such as 2dF. This will enable us to study with high precision the correlation function of these objects over scales in the 2–400 Mpc range, to probe the environmental influence on various type of AGN, and to compare these correlations with those for normal galaxies. The distinct redshift distributions of X-ray selected type I and II AGN can be accurately determined, and this will provide interesting constraints on models of black hole formation compared with models of star formation. The proposed studies will provide a comparison of the clustering properties, as a function of redshift, of X-ray, optical (2dF) or radio selected AGN/QSO.

## 8.4. Survey products

A dedicated `mysql` database with a Java front end interface is available via a site at IASF Milano<sup>37</sup> with a mirror at ESO Santiago. This includes the X-ray source catalogue, and complete catalogues of surveys performed by the LSS Consortium in other wavebands, together with selected subsets from the surveys performed by other Consortia made available under agreement. It also gives access to a selection of data products. The database, currently accessible and used internally by the Consortium, will be gradually opened to the public. The first public release, based on the AO-1 pointings (X-ray source lists and available identifications) is foreseen by the end of 2004.

## Acknowledgments

XMM is an ESA science mission with instruments and contributions directly funded by ESA Member States and NASA. SDS was supported by a post-doctoral position from the Centre National d'Etudes Spatiales. MP and IV are grateful to the ESO/Santiago Office for Science, for a two-week stay in October 2002, where the analysis of the VLT data presented here was initiated; they would like to thank G Marconi for his support at the VLT. Thanks also to G Boese and R Gruber at MPE for enlightening discussions about the RASS PSF.

## References

- [1] Adami C, Mazure A, Katgert P and Biviano A, 1998 *Astron. Astrophys.* **336** 63 [SPIRES]
- [2] Andreon S *et al*, 2004 *Mon. Not. R. Astron. Soc.* **353** 368 [[astro-ph/0405617](#)]
- [3] Andreon S *et al*, 2005 in preparation
- [4] Arnaud M and Evrard A, 1999 *Mon. Not. R. Astron. Soc.* **305** 631
- [5] Baldi A, Molendi S, Comastri A, Fiore F, Matt G and Vignali C, 2002 *Astrophys. J.* **564** 190 [SPIRES]
- [6] Bertin M and Arnouts S, 1996 *Astron. Astrophys. Suppl.* **117** 393
- [7] Böhringer H *et al*, 2000 *Astrophys. J. Suppl.* **129** 435
- [8] Böhringer H *et al*, 2001 *Astron. Astrophys.* **369** 826 [SPIRES]
- [9] Bolzonella M, Miralles J-M and Pelló R, 2000 *Astron. Astrophys.* **363** 476 [SPIRES]
- [10] Browne I W, Mao S, Wilkinson P N, Kus A J, Marecki A and Birkinshaw M, 2000 *SPIE* **4015** 299
- [11] Bruzual G and Charlot S, 1993 *Astrophys. J.* **405** 538 [SPIRES]
- [12] Castander F J *et al*, 1995 *Nature* **281** 59 [SPIRES]
- [13] Cohen A *et al*, 2003 *Astrophys. J.* **591** 640 [SPIRES]
- [14] Collins C A, Burke D J, Romer A K, Sharples R M and Nichol R C, 1997 *Astrophys. J.* **479** L117 [SPIRES]
- [15] de Grandi S *et al*, 1999 *Astrophys. J.* **514** 148 [SPIRES]
- [16] Donahue M, Mack J, Scharf C, Lee P, Postman M, Rosati P, Dickinson M, Voit G M and Stocke J T, 2001 *Astrophys. J.* **552** L93 [SPIRES]
- [17] Ebeling H, Edge A C and Henry J P, 2001 *Astrophys. J.* **553** 668 [SPIRES]
- [18] Ebeling H, Edge A C, Allen S W, Crawford C S, Fabian A C and Huchra J P, 2000 *Mon. Not. R. Astron. Soc.* **318** 333
- [19] Ebeling H, Mullis C R and Tully R B, 2002 *Astrophys. J.* **580** 774 [SPIRES]
- [20] Gioia I M, Henry J P, Maccacaro T, Morris S L, Stocke J T and Wolter A, 1990 *Astrophys. J. Lett.* **356** L35
- [21] Gladders M D and Yee H K, 2000 *Astron. J.* **120** 2148 [SPIRES]
- [22] Haiman Z, Mohr J J and Holder G P, 2001 *Astrophys. J.* **553** 545 [SPIRES]
- [23] Henry J P, Gioia I M, Mullis C R, Voges W, Briel U G and Böhringer H, 2001 *Astrophys. J. Lett.* **553** L109
- [24] Hu W, 2003 *Preprint* [astro-ph/0301416](#)
- [25] Jones L R, Scharf C, Ebeling H, Perlman E, Wegner G, Malkan M and Horner D, 1998 *Astrophys. J.* **495** 100 [SPIRES]

<sup>37</sup> <http://cosmos.mi.iasf.cnr.it/~lssadmin/Website/LSS/>



- [26] Landy S D *et al*, 1996 *Astrophys. J.* **456** L1 [SPIRES]
- [27] Le Fevre O *et al*, 2004 *Astron. Astrophys.* **417** 839 [SPIRES]
- [28] Liang H, 2001 *Preprint astro-ph/0110518*
- [29] Lo K Y, Chiueh T, Liang H, Ma C P, Martin R, Ng K-W, Pen U L and Subramanyan R, 2000 *IAU Symp.* **201** 31
- [30] Lonsdale C *et al*, 2003 *Publ. Astron. Soc. Pac.* **115** 89
- [31] Lobo C, Iovino A, Lazzati D and Chincarini G, 2000 *Astron. Astrophys.* **360** 896 [SPIRES]
- [32] Moretti A, Campana S, Lazzati D and Tagliaferri G, 2003 *Astrophys. J.* **588** 656 [SPIRES]
- [33] Pacaud *et al*, 2005 *A&A*, in preparation
- [34] Panzera M R, Campana S, Covino S, Lazzati D, Mignani R P, Moretti A and Tagliaferri G, 2003 *Astron. Astrophys.* **399** 351 [SPIRES]
- [35] Perlman E S, Horner D J, Jones L R, Scharf C, Ebeling H, Wegner G and Malkan M, 2002 *Astrophys. J. Suppl.* **140** 265
- [36] Peterson J R, Kahn S M, Paerels F B S, Kaastra J S, Tamura T, Bleeker J A M, Ferrigno C and Jernigan J G, 2003 *Astrophys. J.* **590** 207 [SPIRES]
- [37] Press W H and Schechter P, 1974 *Astrophys. J.* **187** 425 [SPIRES]
- [38] Refregier A, Valtchanov I and Pierre M, 2002 *Astron. Astrophys.* **390** 1 [SPIRES]
- [39] Romer A K *et al*, 2000 *Astrophys. J. Suppl.* **126** 206
- [40] Romer A K, Viana P T P, Liddle A R and Mann R G, 2001 *Astrophys. J.* **547** 594 [SPIRES]
- [41] Rosati P, della Ceca R, Norman C and Giacconi R, 1998 *Astrophys. J. Lett.* **492** 21L
- [42] Rosati P, Borgani S and Norman C, 2002 *Ann. Rev. Astron. Astrophys.* **40** 539 [SPIRES]
- [43] Starck J-L and Pierre M, 1998 *Astron. Astrophys.* **128** 397 [SPIRES]
- [44] Valtchanov I, Pierre M and Gastaud R, 2001 *Astron. Astrophys.* **370** 689 [SPIRES]
- [45] Valtchanov I *et al*, 2004 *Astron. Astrophys.* **423** 75 [SPIRES] [[astro-ph/0305192](#)]
- [46] Vikhlinin A, McNamara B R, Forman W, Jones C, Quintana H and Hornstrup A, 1998 *Astrophys. J.* **502** 558 [SPIRES]
- [47] Willis J *et al*, 2004 *MNRAS* submitted
- [48] Xu C K, Lonsdale C J, Shupe D L, Franceschini A, Martin C and Schiminovich D, 2003 *Astrophys. J.* **587** 90 [SPIRES]